Available online at www.sciencedirect.com
SCIENCE DIRECT®
ELSEVIER
Discrete Mathematics 270 (2003) 345–346
DISCRETE MATHEMATICS
www.elsevier.com/locate/disc

Author index to volume 270

Armanious, M.H., Semi-planar Steiner loops of cardinality $2n$ (1–3) 291–298

Bokler, M., Lower bounds for the cardinality of minimal blocking sets in projective spaces (1–3) 13–31
Bonato, A., Homomorphisms and amalgamation (1–3) 33–42
Bonnecaze, A., Y. Choie, S.T. Dougherty and P. Solé, Splitting the shadow (1–3) 43–60
Böttcher, M. and U. Knauer, Postscript: "Endomorphism spectra of graphs" [Discrete Mathematics 109 (1992) 45–57] (1–3) 329–331

Chastand, M. and N. Polat, Reconstruction of infinite locally finite connected graphs (1–3) 61–98
Chen, G.-L., see J.-H. Yin (1–3) 319–327
Choie, Y., see A. Bonnecaze (1–3) 43–60

Deng, D., R. Rees and H. Shen, Further results on nearly Kirkman triple systems with subsystems (1–3) 99–114
Dougherty, S.T., see A. Bonnecaze (1–3) 43–60

Egawa, Y., M. Hagita, K. Kawarabayashi and H. Wang, Covering vertices of a graph by k disjoint cycles (1–3) 115–125

Friedl, K. and L. Rónyai, Order shattering and Wilson's theorem (1–3) 127–136

Hagita, M., see Y. Egawa (1–3) 115–125
Hinrichs, A. and C. Richter, New sets with large Borsuk numbers (1–3) 137–147
Hou, X.-D., Rings and constructions of partial difference sets (1–3) 149–176

Kawarabayashi, K., see Y. Egawa (1–3) 115–125
Knauer, U., see M. Böttcher (1–3) 329–331
Kojima, T., Bandwidth of the composition of two graphs (1–3) 299–309
Kremer, D., Postscript: "Permutations with forbidden subsequences and a generalized Schröder number" [Discrete Mathematics 218 (2000) 121–130] (1–3) 333–334

Lehner, F., Cumulants, lattice paths, and orthogonal polynomials (1–3) 177–191
Lemos, M., Matroids with many common bases (1–3) 193–205
Li, J.-S., see J.-H. Yin (1–3) 319–327

McKee, T.A., Subgraph trees in graph theory (1–3) 3–12
Meng, J. and Z. Zhang, Number of minimum vertex cuts in transitive graphs (1–3) 311–317

Napolitano, V., Erratum to "On a class of finite linear spaces with few lines" [Discrete Mathematics 270 (2003) 207–224] (1–3) 225–226
Napolitano, V., On a class of finite linear spaces with few lines (1–3) 207–224

0012-365X/03/$ - see front matter
doi:10.1016/S0012-365X(03)00335-2

Polat, N., see M. Chastand (1–3) 61–98
Pretzel, O., On reorienting graphs by pushing down maximal vertices—II (1–3) 227–240

Rautenbach, D. and L. Volkmann, On $\alpha,\gamma,(k)$-perfect graphs (1–3) 241–250
Rees, R., see D. Deng (1–3) 99–114
Richter, C., see A. Hinrichs (1–3) 137–147
Rónyai, L., see K. Friedl (1–3) 127–136

Shen, H., see D. Deng (1–3) 99–114
Shi, L., Upper bounds for Ramsey numbers (1–3) 251–265
Solé, P., see A. Bonnecaze (1–3) 43–60
Steinsky, B., Enumeration of labelled chain graphs and labelled essential directed acyclic graphs (1–3) 267–278

Volkmann, L., see D. Rautenbach (1–3) 241–250

Wang, H., see Y. Egawa (1–3) 115–125

Yin, J.-H., J.-S. Li and G.-L. Chen, The smallest degree sum that yields potentially ${}_kC_r$-graphic sequences (1–3) 319–327
Young, P.T., Congruences for degenerate number sequences (1–3) 279–289

Zhang, Z., see J. Meng (1–3) 311–317